

Total number of printed pages-8

3 (Sem-2/CBCS) STA HC2

2022

**STATISTICS**

(Honours)

Paper : STA-HC-2026

(Algebra)

Full Marks : 60

Time : Three hours

**The figures in the margin indicate full marks for the questions.**

1. Answer the following (*any seven*) as directed : 1×7=7
- (a) A polynomial is said to be complete if all the coefficient are present in the polynomial. (State True or False)
- (b) If  $\alpha, \beta, \gamma$  are the roots of the equation  $x^3 - 5x^2 - 16x + 80 = 0$ , then the product of the roots is
- (i) 5
- (ii) -16

Contd.

(iii) -80

(iv) None of the above

*(Choose the correct option)*

(c) State the condition that a matrix  $A$  has to satisfy to be an orthogonal matrix.

(d) A matrix  $A$  will be an Involutary matrix if  $A^2 = \dots\dots\dots$  *(Fill in the blank)*

(e) If two rows or two columns of a determinant be identical, the value of the determinant is

(i) 0

(ii) 1

(iii) None of the above.

*(Choose the correct option)*

(f) If  $A$  be any  $n$ -rowed square matrix, then  $(Adj A) A = A (Adj A) = \dots\dots\dots$

*(Fill in the blank)*

(g) The rank of a unit matrix of order  $n$  is

(i) 1

(ii)  $n$

(iii)  $n-1$

(iv) None of the above

*(Choose the correct option)*

(h) Given that for a  $(5 \times 5)$  matrix  $A$  and  $|A| = 59$ , find  $|3A|$ .

(i) Consider the system of homogeneous linear equations

$$(A)_{m \times n} (X)_{n \times 1} = (O)_{m \times 1}$$

and suppose  $\rho(A) = r$ . Find out the number of linearly independent solutions for this system of equations.

(j) If  $A$  and  $B$  are two equivalence matrices, then  $\text{rank}(A) = \text{rank}(B)$ .

(State True or False)

2. Answer **any four** of the following questions :  
 $2 \times 4 = 8$

(a) Solve the equation  $x^3 - 3x^2 + 4 = 0$ , given that two of its roots being equal.

(b) Examine whether the set

$$S = \left\{ \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}, \begin{pmatrix} 5 \\ 6 \\ 7 \end{pmatrix} \right\} \text{ is}$$

linearly independent or not.

- (c) If  $A$  and  $B$  are symmetric matrices, then show that  $AB$  is symmetric iff  $A$  and  $B$  are commute.
- (d) Show that the necessary and sufficient condition for a square matrix  $A$  to possess the inverse is that  $|A| \neq 0$ .
- (e) Show that if two adjacent rows or columns of a determinant are interchanged, the sign of the determinant is changed, whereas its numerical value remaining the same.
- (f) Given for a  $(3 \times 3)$  matrices  
 $|adj A| = 20$ , find  $|A|$ .
- (g) Write down the matrix of the following forms and verify that it can be written as matrix products  $X'AX$ .
- $$x^2 - 18x_1x_2 + 5x^2$$
- (h) Show that  $\lambda$  is a characteristic root of the matrix  $A$  if and only if there exists a non-zero vector  $X$  such that  $AX = \lambda X$ .

3. Answer **any three** of the following questions : 5×3=15

(a) If  $\alpha, \beta, \gamma$  are the roots of the equation

$x^3 + qx + r = 0$ , find the value of

(i)  $(\beta + \gamma)^{-1} + (\gamma + \alpha)^{-1} + (\alpha + \beta)^{-1}$

(ii)  $\sum (\beta + \gamma - \alpha)^3$

(b) Show that

(i) Every subspace,  $S$ , of  $V_n$  has a basis.

(ii) The row rank of a matrix is the same as its rank.

(c) Prove that

$$\begin{vmatrix} (b+c)^2 & a^2 & a^2 \\ b^2 & (c+a)^2 & b^2 \\ c^2 & c^2 & (a+b)^2 \end{vmatrix} = 2abc(a+b+c)^3$$

(d)  $P, Q$  are non singular matrices. Show

that if  $A = \begin{bmatrix} P & 0 \\ 0 & Q \end{bmatrix}$ , then

$$A^{-1} = \begin{bmatrix} P^{-1} & 0 \\ 0 & Q^{-1} \end{bmatrix}$$

(e) State and prove Cayley-Hamilton theorem.

(f) Show that if  $A$  is an idempotent matrix with dimension  $n \times n$ , then

$$\text{rank}(A) + \text{rank}(I - A) = n$$

(g) Define positive definite, negative definite and semi-positive definite matrices with examples.

4. Answer the following questions (**any three**):  
10×3=30

(a) (i) Derive the standard form of a cubic equation. 5

(ii) Solve the equation by Cardon's method

$$x^3 - 9x - 28 = 0 \quad 5$$

(b) (i) Show that if  $A, B$  are two  $n$ -rowed square matrices then

$$\text{rank}(AB) \geq \text{rank}(A) + \text{rank}(B) - n$$

7

(ii) Show that the vectors

$X_1 = (1, 2, 3)$ ,  $X_2 = (2, -2, 0)$  form a linearly independent set. 3

- (c) Find the inverse of the matrix

$$S = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} \text{ and show that the}$$

transform of the matrix

$$A = \begin{bmatrix} b+c & c+a & b-c \\ c-b & c+b & a-b \\ b-c & a-b & a+b \end{bmatrix} \text{ by } S$$

is a diagonal matrix.

- (d) Show that the equations

$$x + y + z = 6$$

$$x + 2y + 3z = 14$$

$$x + 4y + 7z = 30$$

are consistent and solve them.

- (e) Show that the every  $m \times n$  matrix of rank 'r' can be reduced to the

form  $\begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}$  by a finite chain of

$E$ -operations, where  $I_r$  is the  $r$ -rowed unit matrix.

- (f) Show that a necessary and sufficient condition for a real quadratic form  $X'AX$  to be positive definite is that the leading principal minors of the matrix  $A$  of the form are all positive.

(g) Suppose  $X = \begin{bmatrix} 1 & 2 \\ 0 & -1 \\ 3 & 1 \end{bmatrix}$

Evaluate  $M = I - X(X'X)^{-1}X'$ , where notations have their usual meanings. Show that  $M = M^2$  and find the rank of  $M$  and  $M^2$ .

- (h) Determine the characteristic roots and the corresponding characteristic vectors of the matrix

$$A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$$

---